

**The June 2013 flood in the Upper Danube basin**

G. Blöschl et al.

This discussion paper is/has been under review for the journal Hydrology and Earth System Sciences (HESS). Please refer to the corresponding final paper in HESS if available.

# The June 2013 flood in the Upper Danube basin, and comparisons with the 2002, 1954 and 1899 floods

G. Blöschl, T. Nester, J. Komma, J. Parajka, and R. A. P. Perdigão

Institute of Hydraulic Engineering and Water Resources Management,  
Vienna University of Technology, Vienna, Austria

Received: 5 July 2013 – Accepted: 15 July 2013 – Published: 19 July 2013

Correspondence to: G. Blöschl (bloeschl@hydro.tuwien.ac.at)

Published by Copernicus Publications on behalf of the European Geosciences Union.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



## Abstract

The June 2013 flood in the Upper Danube basin was one of the largest floods in the past two centuries. An atmospheric blocking situation produced precipitation exceeding 300 mm over four days at the northern rim of the Alps. The high precipitation along with high antecedent soil moisture gave rise to extreme flood discharges in a number of tributaries including the Tiroler Ache, Saalach, Salzach and Inn. Runoff coefficients ranged from 0.2 in the Bavarian lowlands to 0.6 in the Alpine areas in Austria. Snowfall at high altitudes (above about 1600 m a.s.l.) reduced the runoff volume produced. Precipitation was distributed over two blocks separated by a few hours which resulted in a single peak, long duration flood wave at the Inn and Danube. At the confluence of the Bavarian Danube and the Inn, the small time lag between the two flood waves exacerbated the downstream flood at the Danube. Because of the long duration and less inundation, there was less flood peak attenuation along the Austrian Danube reach than for the August 2002 flood. Maximum flood discharges of the Danube at Vienna were about  $11\,000\text{ m}^3\text{ s}^{-1}$ , as compared to 10 300, 9600 and  $10\,500\text{ m}^3\text{ s}^{-1}$  in 2002, 1954 and 1899, respectively. This paper reviews the meteorological and hydrological characteristics of the event as compared to the 2002, 1954 and 1899 floods, and discusses the implications for hydrological research and flood risk management.

## 1 Introduction

In June 2013 a major flood struck the Upper Danube basin causing heavy damage along the Danube and numerous tributaries. The city centre of Passau at the confluence of the Danube, Inn and Ilz experienced flood levels that were similar to the highest recorded flood in 1501. Extraordinary flood discharges were recorded along the Saalach and Tiroler Ache at the Austrian–Bavarian border. The flood discharge of the Danube at Vienna exceeded those observed in the past two centuries. First damage estimates suggest figures of several billion Euros (Versicherungsjournal, 2013).

HESSD

10, 9533–9573, 2013

## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

⏪

⏩

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



The June 2013 flood comes at a time with an amazing history of recent large floods. In August 2005, the Danube tributaries in western Tyrol and the south of Bavaria were flooded through extensive precipitation and high antecedent soil moisture (BLU, 2006). In August 2002, a major flood hit the entire Upper Danube basin. Damage was most severe at the northern tributaries of the Austrian Danube at the Czech border, in particular the Aist and Kamp rivers. At the Kamp, flood discharges were almost three times the largest flood in the century before (Gutknecht et al., 2002). Flooding was extensive along the entire Austrian Danube which resulted in the use of the term “century flood”. The preceding decades were relatively flood poor at the Danube with more minor floods in 1991, 1966 and 1965 but a very large flood occurred in July 1954 with major damage along the entire Upper Danube. Again, a couple of decades with almost no floods preceded. The flood of September 1899, then, was the largest measured flood along the Danube with 48 h precipitation totals exceeding 200 mm over an area of 1000 km<sup>2</sup> (Kresser, 1957). Major floods occurred in August 1897, February 1862 and October 1787 with a long record of previous events (Kresser, 1957).

The magnitude of the June 2013 flood has been enormous. The aim of this paper is to analyse the causal factors of this flood including the atmospheric situation, runoff generation and the propagation of the flood wave along the Danube and tributaries. Given the extraordinary nature of the 2013 flood, the paper also compares this flood with the largest Upper Danube floods in the past two centuries, i.e., the floods in August 2002, July 1954 and September 1899.

## 2 The Upper Danube basin

The Upper Danube basin consists of two main subcatchments, the Bavarian Danube and the Inn. The Bavarian Danube catchment in the northwest comprises lowlands with diverse geology. Quaternary and tertiary deposits prevail which are highly permeable and provide large subsurface storage in porous aquifers, and there is also karst in the northwest. Some of the tributaries such as the Lech and Isar originate from the

Alps. Elevations range from 310 to 3000 m.a.s.l. Mean annual precipitation ranges from 650 to more than 2000 mm $\text{yr}^{-1}$  resulting in mean annual runoff depths from 100 to 1500 mm $\text{yr}^{-1}$  (BMU, 2003). The Inn catchment, further in the south, drains a large part of the Austrian Alps. An important tributary is the Salzach. Geologically, the Inn catchment mainly consists of the Northern Calcareous Alps, the Palaeozoic Greywacke zone further in the south and the Crystalline zone along the ridge of the Eastern Alps (Janoschek and Matura, 1980). Elevations range from 310 to 4000 m. Mean annual precipitation ranges from 600 to more than 2000 mm $\text{yr}^{-1}$ , resulting in mean annual runoff depths from 100 to 1600 mm $\text{yr}^{-1}$  (Parajka et al., 2007; Nester et al., 2011).

The Bavarian Danube and the Inn join at Passau. Downstream of the confluence, along the Austrian reach of the Danube, southern tributaries from the high rainfall areas in the Calcareous Alps include the Traun, Enns and Ybbs. The northern tributaries from the lower rainfall areas with mainly granitic geology include the Aist and Kamp. Flood protection levees have been built along many tributaries and the Danube itself during the 19th and 20th century. The total catchment area of the Danube at Wildungsmauer downstream of Vienna is 104 000 km<sup>2</sup>. Figure 1 shows the catchments discussed in this paper, and Table A1 gives their main characteristics.

### 3 Large scale atmospheric conditions

The 2013 flood was produced by an atmospheric situation that is typical of floods in the Upper Danube. In the second half of May 2013 the planetary waves of the large-scale atmospheric flow regime in the Northern Hemisphere exhibited stationary behaviour as the mean eastward zonal flow decelerated and no longer exceeded the westward propagation of the Rossby waves produced by the latitude-varying Coriolis effect. The stationary nature of the systems is illustrated by the five-day moving averages of the geopotential height centred on 26 May and 30 May 2013 in Fig. 2. The shapes of the system centres in Fig. 2 are circular or near-circular indicating that these centres have barely moved during the five-day averaging periods. The persistence is further

## HESSD

10, 9533–9573, 2013

### The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion







The end-of-May soil moisture exceeded all values for this time of the year over the period 1962 to 2013 considered in the simulations of BfG (2013). Also, ground water levels were particularly high. Further in the south, the soils were still relatively wet although the situation was less extreme. Lofer (near the Weißbach gauge in Fig. 1), for example, had 209 mm of precipitation in May 2013 as compared to the long term May mean of 140 mm. This means that, at the beginning of the event the soils were wet throughout the Upper Danube basin, although there was a pronounced north–south gradient with higher soil moisture in the north, and lower soil moisture in the south.

Heavy precipitation started on 29 May in the northern part of the Bavarian Danube catchment around the city of Regensburg. From 29 May to 4 June precipitation totals of 95 mm were observed in Regensburg, 94 mm in Straubing, and 112 mm at Großer Arber to the east of Regensburg. In the southern part of the Bavarian Danube catchment and the Inn catchment, heavy precipitation started on 30 May and lasted until 2 June 2013, with smaller intensities on 3 and 4 June. From 29 May to 4 June precipitation totals of 232 mm were observed in Lofer, 244 mm in Kössen (25 km northwest of Lofer) and 270 mm in Samerberg (40 km northwest of Lofer). Figure 4 (top left panel) shows the spatial pattern of precipitation for a period of seven days (29 May to 4 June 2013). As indicated in the figure, precipitation was highest along the northern ridge of the Alps in Austria (Tirol, Salzburg and Upper Austria) and there was also very significant precipitation further in the north. Precipitation interpolated between the rain gauges based on weather radar exceeded 300 mm during this time period (Fig. 4).

As an example, Fig. 5 shows the evolution of catchment precipitation during May and the first days of June 2013 for the Weißbach catchment, a tributary to the Salzach (Fig. 1). Catchment precipitation was estimated from the gridded data as in Fig. 4. In this catchment, May 2013 precipitation was 184 mm, and the event precipitation from 30 May to 2 June was an additional 175 mm. The event consisted of two main precipitation blocks separated by a few hours of no or lower intensity rain. These two rain blocks were apparent over most of the Upper Danube basin.

## HESSD

10, 9533–9573, 2013

### The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion







(Kresser, 1957). Snow also played a role. Snow accumulated above about 1400 m which did not contribute to the event. Most of the precipitation fell in the Inn, Salzach, Traun and Enns catchments. There was a much smaller contribution from the Bavarian Danube (Lauda, 1900).

## 5 Runoff generation

### 5.1 Runoff generation of the 2013 flood

Table 1 gives the event characteristics of a number of catchments with particularly high event precipitation. The catchments drain into the Inn (Rosenheim, St. Johann, Staudach), Salzach (Weißbach, Obergäu) or directly into the Austrian Danube (Wels) (see Fig. 1). The event precipitation was of the same order of magnitude as the antecedent precipitation in May, ranging from 160 mm in the Obergäu catchment to 231 mm in the Rosenheim catchment. Because of the relatively high antecedent precipitation, and hence soil moisture, the runoff coefficients are quite large. They ranged between 0.42 and 0.56. However, these runoff coefficients are not unusually high. For comparison, the average runoff coefficients in these catchments for events with more than 50 mm precipitation are given. In Obergäu and Wels, the runoff coefficients of the 2013 event are 0.58 and 0.56 and thus very similar to the average runoff coefficients of all large events (precipitation > 50 mm) in these catchments. In St. Johann, Staudach and Weißbach the runoff coefficients of the 2013 event are 0.42 which is somewhat lower than the averages of the large events, in spite of the relatively wet antecedent conditions. This is because part of the precipitation fell as snow and remained as snow cover until after the event in the highest parts of the catchment.

Figure 6 (left panel) shows the 2013 event for the Weißbach catchment in more detail. The cumulative precipitation illustrates the two precipitation blocks. The first block of 45 mm started on 30 May around midday and lasted until the evening of 31 May. The second block of 130 mm started on 1 June in the afternoon and lasted until the

## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



evening of 2 June. The first block of precipitation led to a slight increase in runoff, while the second block of precipitation increased the event runoff to a total of 74 mm. There are two reasons for the very small response to the first precipitation block. The first is the soil storage left at the beginning of the event. The second are the temperatures which were below 0 °C in one third of the catchment (light green line in Fig. 6) so there was significant snow fall. During the second block of precipitation the percentage of the catchment with temperatures below 0 °C was lower, particularly at the beginning of the second block when it was warmer. The figure suggests that, over the entire event, around 25 % of the precipitation, or 44 mm, fell as snow, the remaining 131 mm fell as rain. If one only counts rainfall, the associated runoff coefficient is 0.56 which is more in line with the antecedent soil moisture. However, it is likely that some (but not all) of the snow that fell at the beginning of the event melted during the event. The situation in the Weißbach is typical of the Alpine high rainfall catchments during the June 2013 flood. St. Johann and Staudach (Table 1) give similar figures but somewhat higher runoff coefficients because of the lower fraction of snow fall.

In the Bavarian Danube catchment there was also substantial precipitation albeit with a strong south-north gradient and precipitation started earlier. Figure 6 (right panel) shows the cumulative event precipitation and event runoff for the Hofkirchen catchment (45 610 km<sup>2</sup>). Catchment precipitation was significantly lower than in the Weißbach with a total of 116 mm since the northwest of the catchment contributed less precipitation than the southern, Alpine part of it. However, relative to previous events, the precipitation in this catchment was still enormous. At the scale of this catchment, the two precipitation blocks are still visible, but they were only separated by 12 h. The first block consisted of 45 mm, the second of 61 mm, and there was some early precipitation of 10 mm on 29 May. Temperatures were above zero in almost the entire catchment. Only 31 mm out of the 116 mm precipitation contributed to the runoff resulting in a runoff coefficient of 0.27. This is because of the highly permeable soils and the large storage capacity in the catchment in spite of the high antecedent soil moisture. The cumulative runoff depth in Fig. 6 (right panel, red line) indicates the substantial delay between

## HESSD

10, 9533–9573, 2013

### The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



precipitation and runoff. Essentially all the precipitation was stored in the catchment at the inception of runoff.

## 5.2 Comparison with previous floods

It is now of interest to put the 2013 event where rainfall was most severe into the context of the longer term flood history in the area. Figure 7 (left panel) shows the extreme value statistics for 48 h precipitation of the Lofer rain gauge located near the Weißbach stream gauge. In June 2013, the two-day precipitation was the highest since beginning of the data set in 1961. Around 173 mm of precipitation were observed in two days at this station, as compared to 122 mm during the August 2002 flood. Based on a Gumbel distribution, this precipitation total corresponds to a return period of about 70 yr. The right panel of Fig. 7 shows the extreme value statistics of the flood peaks at the Weißbach stream gauge. With a peak runoff of  $480 \text{ m}^3 \text{ s}^{-1}$  the 2013 flood was the highest since beginning of the observations in 1959. The associated return period is around 100 yr. The 2002 flood peak was slightly lower with a peak discharge of  $400 \text{ m}^3 \text{ s}^{-1}$ . The return period on the order of 100 yr for the June 2013 flood peak applies to a number of catchments in the area, in particular those at the Tiroler Achen und Saalach (Table 1, Fig. 1), and produced flood discharges with return periods of about 100 yr at the downstream reaches along the Salzach and Inn.

At the small catchment scale, the August 2002 flood was most severe in the Kamp catchment at the Austrian–Czech border. In the Kamp and nearby catchments, the soils are sandy and thus much more permeable than the catchments of Table 1. Because of this, the runoff coefficients are usually quite low. For the Kamp at Zwettl catchment ( $620 \text{ km}^2$ ) the average runoff coefficient (for event precipitation  $> 50 \text{ mm}$ ) is 0.20 (Merz et al., 2006). Due to the large rainfall depths of the 2002 event (200 and 115 mm from the two events) soils did become saturated leading to runoff coefficients of 0.41 and 0.58 for the two events, respectively (Komma et al., 2007) which is almost three times the average. The resulting flood peak was  $460 \text{ m}^3 \text{ s}^{-1}$  (as compared to the second largest peak of  $170 \text{ m}^3 \text{ s}^{-1}$  since beginning of the record in 1896) which made the

## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion





The Inn exhibited a much faster response as is always the case with this type of regional floods. The Upper Inn showed very little flood runoff and the flood wave built up through tributaries in Bavaria. The flood wave of the Inn at Wasserburg merged with the Salzach wave, peaking essentially at the same time, and produced a very steep wave at Schärding. There was relatively little flood retention along the Inn before the confluence but a number of small tributaries such as the Rott ( $280 \text{ m}^3 \text{ s}^{-1}$  peak flow associated with a return period around 50 yr) contributed to the magnitude of the flood. At Schärding the flood peaked at 15:00 LT on 3 June with a discharge of about  $5950 \text{ m}^3 \text{ s}^{-1}$  which represents an estimated return period of about 100 yr.

The confluence of the Inn with the Bavarian Danube at Passau resulted in a characteristic, combined shape of the flood wave at Achleiten where the fast and slow contributions of Inn and Danube are clearly visible. During the propagation of the flood wave along the Austrian Danube, it changed shape due to retention in the flood plains, which is apparent by the kink of the rising limb about a day before the peak. Inflow from southern tributaries along the Austrian reach of the Danube, including the Traun, Enns and Ybbs, gave rise to an early secondary peak, indicating that these tributaries peaked much earlier and hardly contributed to peak flows along the Danube. While the flood peaked on 2 June at 20:00 LT in both Wasserburg and Oberndorf, it peaked on 6 June at 05:00 LT in Wildungsmauer close to the Austrian–Slovak border.

## 6.2 Confluence at Passau of the 2013 flood and comparison with the 2002, 1954, 1899 floods

The confluence of the peaky flood wave from the Inn with the more delayed flood wave from the Bavarian Danube at Passau is a crucial element in the flood characteristics of the Upper Danube. During the 2013 flood, the inundation level in Passau was enormous (12.89 m) (BfG, 2013). It was of the same order of magnitude as the 1501 flood event (between 12.70 and 13.20 m, depending on the source) which is considered the highest flood in the past millennium (Schmidt, 2000; BfG, 2013).

# HESSD

10, 9533–9573, 2013

## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



To understand the particularly large flood levels in Passau with implications downstream Fig. 9 shows the flood hydrographs for stream gauges at the Inn and Danube just above the confluence (Schärding and Hofkirchen/Vilshofen), as well as just below the confluence (Achleiten). For the 2013 and 2002 floods the discharges are given, while for the 1954 and 1899 floods the water levels are given since reconstruction of the complete runoff hydrograph is prone to uncertainties. It is clear that the Inn flood wave is always much faster than that from the Bavarian Danube. In 2013, the flood wave of the Bavarian Danube (at Hofkirchen) arrived somewhat earlier than usual, as compared to the Inn flood wave. This is because rainfall started a little earlier, on 29 May, rather than on 30 May as in the Inn and because of the very high antecedent soil moisture. On 4 June 2013, a dam along the Bavarian Danube and the Isar collapsed and caused large spatial flooding in Deggendorf, 30 km upstream of Hofkirchen. This explains the kink in the runoff hydrograph and the flat crest of the flood wave.

In terms of the confluence, the June 2013 flood was most similar to the July 1954 flood when the Bavarian Danube had similarly large flood discharges. At Hofkirchen (and the nearby Vishofen gauge) the 2013 and 1954 peaks were  $3420$  and  $3320 \text{ m}^3 \text{ s}^{-1}$ , respectively. The 2013 peaks were slightly larger because of three main reasons. (i) In 2013 it was warmer than in 1954 with less snowfall, so more rain was available for runoff generation; (ii) in 2013 rainfall was more concentrated along the northern fringe of the Alps where infiltration capacity is lower than in the north of the basin where much of the rain fell in 1954; (iii) antecedent soil moisture in the north was probably higher in 2013 than in 1954.

During the 2013 flood event, the wave from the Bavarian Danube arrived comparatively early. At the time Schärding peaked, Hofkirchen showed a discharge of  $3000 \text{ m}^3 \text{ s}^{-1}$  which was close to its peak discharge of  $3420 \text{ m}^3 \text{ s}^{-1}$ . With smaller differences in the time lag between the Bavarian Danube and Inn waves and larger discharges in the Bavarian Danube, the resulting flood wave was significantly higher than in 1954 ( $10\,000 \text{ m}^3 \text{ s}^{-1}$  peak flow in Achleiten in 2013 as compared to  $9100 \text{ m}^3 \text{ s}^{-1}$  in 1954).

2002 was different in that the Bavarian Danube flood wave was more delayed, so the superposition was less efficient than in 2013 and 1954. The Inn wave was smaller too because of less rainfall in the catchment area. The 2002 flood was therefore much smaller at Passau, but received major contributions from the tributaries along the Austrian reach of the Danube (including the Kamp, Aist, Traun and Enns) as illustrated by Fig. 10.1 in Blöschl et al. (2013). Similarly, the 1899 flood was much smaller at the Bavarian Danube because of less rainfall and dry soils. The 1899 flood was most severe along the Inn, the Traun and the Enns as indicated by the rainfall distribution in Fig. 3.

### 6.3 Propagation of the 2013 flood along the Austrian Danube and comparison with 2002, 1954, 1899

After the confluence of the Bavarian Danube and the Inn at Passau, the 2013 flood wave travelled downstream the Austrian Danube changing shape and shifting the timing. Figure 10 shows the time evolution of the peak flow from Passau to Wildungsmauer close to the Austrian–Slovak border. For this reach, the total time lag of the peak during the 2013 flood was 60 h. The celerity of the flood wave is related to two main factors, (i) the rainfall distribution and hence the contribution and timing of the tributaries, (ii) the flood propagation of the main wave itself and the associated retention in the flood plains. The propagation of the 1954 flood was much slower than that of the other events (118 h, Fig. 10). This is because much of the rainfall occurred in the Bavarian Danube catchment (Fig. 3) with relatively minor contributions from the southern tributaries Traun and Enns, and hardly any contributions from Kamp and Ybbs. Also, there were significant inundations in the flood plains, in particularly upstream and downstream of Linz, contributing to the delay. In contrast, 2002 was the fastest event which was because those tributaries contributed significantly to the rising limb and peak of the flood wave, thus apparently accelerating the flood propagation, in particular the first wave (2002-1). The 2013 flood was in between 1954 and 1899 in terms of rainfall distribution, so from that perspective one would expect celerities between 1954

HESSD

10, 9533–9573, 2013

## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



and 1899, i.e., a total time lag of around 80 to 90 h. However, the wave celerity was somewhat faster than that. During the 1899 flood, a total area of 1070 km<sup>2</sup> was inundated along the Austrian Danube and tributaries as estimated by Lauda (1900). With an assumed average inundation depth of 2 m, this area corresponds to a total retention volume of more than 2 billion m<sup>3</sup>. This is very significant relative to the volume of the 1899 flood (6.6 billion m<sup>3</sup>, without base flow, Lauda, 1990), indicating that the retention effect was indeed very important during that event. As compared to 1899, the retention volume had been decreased by 2013 because of flood protection levees, hydropower projects and mobile flood protection. The smaller retention volumes, and possibly the greater water depths in the reservoirs, have likely contributed to accelerating the flood wave, although detailed hydrodynamic studies would be needed to exactly ascertain the reasons for the changes in the wave celerities (e.g. Fischer-Antze et al., 2008). Detailed analyses of observed flood celerities at the Danube are provided in Szolgay and Danáčová (2007) and Mitkova et al. (2005).

#### 6.4 Magnitude of the 2013 flood at the Austrian Danube and comparison with 2002, 1954, 1899

Figure 11 shows the extreme value statistics of the maximum annual floods for two stream gauges along the Austrian Danube, the Kienstock and Korneuburg gauges. Korneuburg is about 12 km upstream of Vienna, while Kienstock is about 85 km upstream of Vienna. Even though the spatial distribution of precipitation, runoff generation and the spatial evolution of the 2013 and 2002 floods were different, the maximum runoff values along the Austrian Danube were quite similar. The flood runoff of both events was associated with return periods on the order of 100 yr, based on the statistical analysis. The 2013 flood runoff was clearly larger than that of the 1899 and 1954 floods.

It is interesting to compare the relative magnitudes of the peak runoff of these events between the two locations. While at Kienstock the 2013 flood runoff peak was almost identical to that of 2002, at Korneuburg it was significantly lower. In 2013, the peak runoff only decreased slightly along this reach (Kienstock 11 100 m<sup>3</sup> s<sup>-1</sup>, Korneuburg

## HESSD

10, 9533–9573, 2013

### The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



11 055 m<sup>3</sup> s<sup>-1</sup>). In 2002, this decrease was much stronger (11 300 vs. 10 250 m<sup>3</sup> s<sup>-1</sup>). At first sight this is counterintuitive, as the 2002 flood had substantial inflows along the Kienstock–Korneuburg reach (in particular from the Kamp, Figs. 1 and 3) while the inflows during the 2013 event were much smaller. The smaller peak attenuation is due to two main reasons. (i) Both events, 2002 and 2013, had two precipitation blocks. However, in 2013 the time period between the blocks was only a few hours, while it was four days in 2002. The 2002 event had two atmospheric rainfall producing situations with movement in between due to less atmospheric stationarity than in 2013. Because of the short interstorm period, the two precipitation blocks in 2013 combined to a single flood wave on the Danube which exhibited a larger runoff volume than either of the two 2002 events. This is illustrated in Fig. 12 by the flood hydrographs of the 2002 and 2013 flood events at Kienstock and Korneuburg. During the 2013 flood at Kienstock, a runoff threshold of 9000 m<sup>3</sup> s<sup>-1</sup> was exceeded during 81 h, while in 2002 the same discharge was only exceeded 52 h, so the flood volume to be stored in the flood plains was smaller in 2002 resulting in a bigger peak reduction. (ii) Second, there were differences in the performance of the levees and the way the hydraulic structures were operated along the Danube. In the aftermath of the 2002 flood, levees had been strengthened which resulted in relatively less inundation along the reach in 2013 as compared to 2002 which may have contributed to a smaller peak reduction. Again, detailed hydrodynamic studies would be needed to analyse the effects of individual hydraulic structures.

Overall, the flood volumes at Korneuburg (near Vienna) were enormous for all four floods. The total runoff volume of the 2013 flood from 31 May to 17 June (including base flow) was 9.5 billion m<sup>3</sup>, the volume of the 2002 flood from 11 to 26 August (second event only) was 6.5 billion m<sup>3</sup>, and the volumes of the 1954 and 1899 floods (8–27 July and 10–28 September) were 9.9 and 8.5 billion m<sup>3</sup>, respectively.

## 7 Lessons learnt and implications

### 7.1 Implication for hydrological research and design flood estimation

The June 2013 flood came at a time when the 2002 flood has still been fresh in the minds of hydrologists. The lessons learnt from the 2013 flood therefore complement those learnt from the 2002 flood. From a hydrological perspective, the analysis of the causal factors in this paper highlights the outstanding role of the *combination* of extreme factors that control the magnitude and characteristics of extreme floods. Indeed, the 2013 flood has had a number of characteristics that contributed to increasing its magnitude, in particular relatively high antecedent soil moisture, little shift between the flood peaks at the confluence of the Bavarian Danube and the Inn, and rainfall blocks close together resulting in a single, large volume flood wave with relatively small peak attenuation. While the important role of the combination of a number of extreme factors has been already emphasised in the past (Kresser, 1957; Gutknecht, 1994), the combination of factors remains an essential concept for understanding the magnitude of large, regional floods. On the other hand, there were also factors that could have easily been more extreme. There was less total precipitation than in 1899 and, in the Alpine area, there was a significant snow fall component which retained some of the water in the catchments. Clearly, more extreme situations are possible from a hydrological perspective, albeit unlikely. Rainfall as in 1899 with high antecedent soil moisture like in 2013 would produce a significantly larger flood. This has important implications for estimating design flood values. While statistical analyses are important when estimating design flood discharges, particularly in large catchments with long flood records, it is equally important to address the problem from a process perspective and understand what combinations of factors could plausibly be expected to occur during extreme situations. This is in the spirit of flood frequency hydrology, a framework for understanding and estimating flood discharges by combining local flood data with additional types of information: temporal information on historic floods, spatial information on floods in neighbouring catchments, and causal information on the flood processes (Merz and

## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)





of these effects. In the FloodChange project these changes are analysed in detail on the basis of the flood frequency hydrology concept of combining local flood data with historic floods, and spatial and causal flood information (Merz and Blöschl, 2008a).

## 7.2 Implications for flood risk management

5 While there has been significant damage during the 2013 flood there is no doubt that, overall, the flood protection measures put into place in the past two centuries have vastly reduced the damage relative to a scenario without protection. Historically, much of the mitigation activities have been focused on structural measures such as levees, but there is an increasing awareness that a number of flood management measures are  
10 needed to complement each other (Merz et al., 2010). These include structural measures such as levees for flood protection and construction of polders for flood retention, and non-structural measures such as spatial planning and increasing the preparedness of local citizens. Retaining water in polders and retention basins is always useful as, even for extreme flooding, flood attenuation will occur with positive effects downstream. The drawback is that a lot of area is needed for flood retention to be effective at large  
15 rivers such as the Danube as the flood peak reduction is a direct function of the available storage volume relative of the flood volume. In highly populated areas it is difficult to make sufficiently large areas available, so levees will continue to play a central role in flood management. However, levees may exacerbate flood risk downstream. Integrated  
20 flood risk management therefore considers the river basin as a whole as stipulated by the EU flood risk directive (EU, 2007).

Local protection of buildings along with raising flood risk awareness and preparedness of local citizens may be highly effective to complement the other measures. For these, and other flood event management measures such as early evacuation, reliable warning systems, driven by hydrological forecast models, are needed. While large  
25 scale meteorological models and satellites provide important inputs, in particular on future precipitation, capturing the local hydrological situation is essential for accurately modelling floods (Blöschl, 2008). Increasingly longer lead times are expected from

### The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)





For such extremes, as with all floods, the hallmark of integrated flood risk management is the interplay of all measures in a seamless way.

*Acknowledgements.* We would like to thank all institutions who have provided data, in particular the Hydrologic Offices and the Central Institute for Meteorology and Geodynamics. All data of the 2013 flood are tentative. We would also like to thank the ERC (Advanced Grant on Flood Change) and the FWF (project no P 23723-N21) for financial support.

## References

- BfG: Das Juni-Hochwasser des Jahres 2013 in Deutschland (The 2013 June flood in Germany), BfG Report no. 1793, Federal Institute of Hydrology, Koblenz, Germany, 2013.
- Blöschl, G.: Flood warning – on the value of local information, *International Journal of River Basin Management*, 6, 41–50, 2008.
- Blöschl, G. and Montanari, A.: Climate change impacts-throwing the dice?, *Hydrol. Process.*, 24, 374–381, 2010.
- Blöschl, G., Ardoin-Bardin, S., Bonell, M., Dorninger, M., Goodrich, D., Gutknecht, D., Matorros, D., Merz, B., Shand, P., and Szolgay, J.: At what scales do climate variability and land cover change impact on flooding and low flows?, *Hydrol. Process.*, 21, 1241–1247, 2007.
- Blöschl, G., Sivapalan, M., Wagener, T., Viglione, A., and Savenije, H. H. G. (Eds): *Runoff Prediction in Ungauged Basins – Synthesis across Processes, Places and Scales*, Cambridge University Press, Cambridge, UK, 465 pp., 2013a.
- Blöschl, G., Viglione, A., and Montanari, A.: Emerging approaches to hydrological risk management in a changing world, in: *Climate Vulnerability: Understanding and Addressing Threats to Essential Resources*, Elsevier Inc., Academic Press, 3–10, 2013b.
- BLU (Bayerisches Landesamt für Umwelt): August – Hochwasser 2005 in Südbayern (Aug 2005 flood in Southern Bavaria), Endbericht vom 12.04.2006, Bayerisches Landesamt für Umwelt, München, 49 pp., 2006.
- BMU: *Hydrologischer Atlas von Deutschland (Hydrological Atlas of Germany)*, Bundesministerium für Umwelt, Naturschutz und Reaktorsicherheit, Koblenz, 2003.
- Carr, G., Blöschl, G., and Loucks, D. P.: Evaluating participation in water resource management: a review, *Water Resour. Res.*, 48, W11401, doi:10.1029/2011WR011662, 2012.

## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



Deutsche Vereinigung für Wasserwirtschaft, Abwasser und Abfall (DWA): Merkblatt Ermittlung von Hochwasserwahrscheinlichkeiten (Guidelines for estimating flood probabilities), DWA-M. 552, Hennef, Germany, 2012.

Di Baldassarre, G., Viglione, A., Carr, G., Kuil, L., Salinas, J. L., and Blöschl, G.: Socio-hydrology: conceptualising human-flood interactions, *Hydrol. Earth Syst. Sci. Discuss.*, 10, 4515–4536, doi:10.5194/hessd-10-4515-2013, 2013.

EU: The European Parliament and the Council of the European Union, Directive 2007/60/EC of the European Parliament and the Council of 23 October 2007 on the assessment and management of flood risks, *Off. J. Eur. Union*, L288/27–L288/34, 2007.

Fischer-Antze, T., Olsen, N. R. B., and Gutknecht, D.: Three-dimensional CFD modeling of morphological bed changes in the Danube River, *Water Resour. Res.*, 44, W09422, doi:10.1029/2007WR006402, 2008.

Gaál, L., Szolgay, J., Kohnová, S., Parajka, J., Merz, R., Viglione, A., and Blöschl, G.: Flood timescales: understanding the interplay of climate and catchment processes through comparative hydrology, *Water Resour. Res.*, 48, W04511, doi:10.1029/2011WR011509, 2012.

Godina, R., Lalk, P., Lorenz, P., Müller, G., and Weilguni, V.: Die Hochwasserereignisse im Jahr 2002 in Österreich (The flood events of 2002 in Austria), *Mitt. Hydrogr. Dienstes Österreich*, 82, 1–39, 2003.

Gutknecht, D.: Extremhochwässer in kleinen Einzugsgebieten (Extreme floods in small catchments), *Österreichische Wasser- und Abfallwirtschaft*, 46, 50–57, 1994.

Gutknecht, D., Reszler, C., and Blöschl, G.: Das Katastrophenhochwasser vom 7. Aug 2002 am Kamp – eine erste Einschätzung (The Aug 7, 2002 – flood of the Kamp – a first assessment) *Elektrotechnik und Informationstechnik*, 119, 411–413, 2002.

Haiden, T., Kann, A., Wittmann, C., Pistotnik, G., Bica, B., and Gruber, C.: The integrated now-casting through comprehensive analysis (INCA) system and its validation over the eastern Alpine region, *Weather Forecast.*, 26, 166–183, 2011.

HZB: Das Juli-Hochwasser 1954 im österreichischen Donauegebiet (The July 1954 flood in the Austrian Danube basin), *Beiträge zur Hydrographie Österreichs*, Nr. 29, Hydrogr. Zentralbüro Wien, 139 pp., 1955.

Janoschek, W. R. and Matura, A.: Outline of the Geology of Austria, *Abh. Geol. B.-A.*, 26e, 7–98, 1980.

Kistler, R., Kalnay, E., Collins, W., Saha, S., White, G., Woollen, J., Chelliah, M., Ebisuzaki, W., Kanamitsu, M., Kousky, V., van den Dool, H., Jenne, R., and Fiorino, M.: The NCEP-NCAR

## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[◀](#)

[▶](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)

[Printer-friendly Version](#)

[Interactive Discussion](#)



50 year reanalysis: monthly means CD-ROM and documentation, B. Am. Meteorol. Soc., 82, 247–268, 2001.

Komma, J., Reszler, C., Blöschl, G., and Haiden, T.: Ensemble prediction of floods – catchment non-linearity and forecast probabilities, Nat. Hazards Earth Syst. Sci., 7, 431–444, doi:10.5194/nhess-7-431-2007, 2007.

Komma, J., Blöschl, G., and Reszler, C.: Soil moisture updating by Ensemble Kalman Filtering in real-time flood forecasting, J. Hydrol., 357, 228–242, 2008.

Kresser, W.: Der Einfluß der Regulierungs- und Kraftwerksbauten auf die Hochwasserverhältnisse der österreichische Donau (Effect of river training and power plants on the floods of the Austrian Danube), Oesterreichische Wasserwirtschaft, 6, 65–68, 1954.

Kresser, W.: Die Hochwässer der Donau (The floods of the Danube), Schriftenreihe des österreichischen Wasserwirtschaftsverbandes, Wien, 32–33, 1957.

Kundzewicz, Z. W. (Ed.): Changes in Flood Risk in Europe, IAHS Special Publication 10, IAHS Press, Wallingford, 516 pp., 2012.

Lauda, E.: Die Hochwasserkatastrophe des Jahres 1899 im österreichischen Donaugebiete (The flood disaster of 1899 in the Austrian Danube basin), Beiträge zur Hydrographie Österreichs, IV. Heft, k.k. hydrographisches Central-Bureau, Wien, 1900.

Merz, B., Kreibich, H., Schwarze, R., and Thieken, A.: Review article “Assessment of economic flood damage”, Nat. Hazards Earth Syst. Sci., 10, 1697–1724, doi:10.5194/nhess-10-1697-2010, 2010.

Merz, B., Vorogushyn, S., Uhlemann, S., Delgado, J., and Hündecha, Y.: HESS Opinions “More efforts and scientific rigour are needed to attribute trends in flood time series”, Hydrol. Earth Syst. Sci., 16, 1379–1387, doi:10.5194/hess-16-1379-2012, 2012.

Merz, R. and Blöschl, G.: Flood frequency hydrology: 1. Temporal, spatial, and causal expansion of information, Water Resour. Res., 44, W08432, doi:10.1029/2007WR006744, 2008a.

Merz, R. and Blöschl, G.: Flood frequency hydrology: 2. Combining data evidence, Water Resour. Res., 44, W08433, doi:10.1029/2007WR006745, 2008b.

Merz, R. and Blöschl, G.: A regional analysis of event runoff coefficients with respect to climate and catchment characteristics in Austria, Water Resour. Res., 45, W01405, doi:10.1029/2008WR007163, 2009.

Merz, R., Blöschl, G., and Parajka, J.: Spatio-temporal variability of event runoff coefficients in Austria, J. Hydrol., 331, 591–604, 2006.

## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion

- Merz, R., Parajka, J., and Blöschl, G.: Time stability of catchment model parameters: implications for climate impact analyses, *Water Resour. Res.*, 47, W02531, doi:10.1029/2010WR009505, 2011.
- 5 Mitkova, V., Pekarova, P., Miklanek, P., and Pekar, J.: Analysis of flood propagation changes in the Kienstock–Bratislava reach of the Danube River, *Hydrolog. Sci. J.*, 50, 655–668, 2005.
- Nester, T., Kirnbauer, R., Gutknecht, D., and Blöschl, G.: Climate and catchment controls on the performance of regional flood simulations, *J. Hydrol.*, 402, 340–356, 2011.
- Nester, T., Komma, J., Viglione, A., and Blöschl, G.: Flood forecast errors and ensemble spread – a case study, *Water Resour. Res.*, 48, W10502, doi:10.1029/2011WR011649, 2012a.
- 10 Nester, T., Kirnbauer, R., Parajka, J., and Blöschl, G.: Evaluating the snow component of a flood forecasting model, *Hydrol. Res.*, 43, 762–779, 2012b.
- Parajka, J., Merz, R., and Blöschl, G.: Uncertainty and multiple objective calibration in regional water balance modeling – case study in 320 Austrian catchments, *Hydrol. Process.*, 21, 435–446, 2007.
- 15 Parajka, J., Kohnová, S., Merz, R., Szolgay, J., Hlavcová, K., and Blöschl, G.: Comparative analysis of the seasonality of hydrological characteristics in Slovakia and Austria, *Hydrolog. Sci. J.*, 54, 456–473, 2009.
- Parajka, J., Kohnová, S., Bálint, G., Barbuc, M., Borga, M., Claps, P., Cheval, S., Dumitrescu, A., Gaume, E., Hlavcová, K., Merz, R., Pfaundler, M., Stancalie, G., Szolgay, J., and Blöschl, G.: Seasonal characteristics of flood regimes across the Alpine–Carpathian range, *J. Hydrol.*, 394, 78–89, 2010.
- 20 Peel, M. C. and Blöschl, G.: Hydrologic modelling in a changing world, *Prog. Phys. Geog.*, 35, 249–261, 2011.
- Petrow, T., Zimmer, J., and Merz, B.: Changes in the flood hazard in Germany through changing frequency and persistence of circulation patterns, *Nat. Hazards Earth Syst. Sci.*, 9, 1409–1423, doi:10.5194/nhess-9-1409-2009, 2009.
- 25 Ribicic, H., Waser, J., Fuchs, R., Blöschl, G., and Gröller, E.: Visual analysis and steering of flooding simulations, *IEEE T. Vis. Comput. Gr.*, 19 1062–1075, 2013.
- Salazar, S., Francés, F., Komma, J., Blume, T., Francke, T., Bronstert, A., and Blöschl, G.: A comparative analysis of the effectiveness of flood management measures based on the concept of “retaining water in the landscape” in different European hydro-climatic regions, *Nat. Hazards Earth Syst. Sci.*, 12, 3287–3306, doi:10.5194/nhess-12-3287-2012, 2012.
- 30

- Schmidt, M. (Ed.): Hochwasser und Hochwasserschutz in Deutschland vor 1850 (Floods and flood protection in Germany before 1850), Oldenbourg Industieverlag, Munich, 280 pp., 2000.
- Sivapalan, M., Savenije, H. H. G., and Blöschl, G.: Socio-hydrology: a new science of people and water, *Hydrol. Process.*, 26, 1270–1276, 2012.
- Szolgay, J. and Danáčová, M.: Detection of changes in the flood celerity by multilinear routing on the Danube, *Meteorological Journal*, 10, 219–224, 2007.
- Szolgayova, E., Laaha, G., Blöschl, G., and Bucher, C.: Factors influencing long range dependence in streamflow of European rivers, *Hydrol. Process.*, online first, doi:10.1002/hyp.9694, 2013.
- Ulbrich, U., Brücher, T., Fink, A. H., Leckebusch, G. C., Krüger, A., and Pinto, J. G.: The central European floods of Aug 2002: Part 1 – Rainfall periods and flood development, *Weather*, 58, 371–377, 2003.
- Versicherungsjournal: available at: <http://www.versicherungsjournal.at/markt-und-politik/hochwasser-macht-frage-der-versicherbarkeit-virulent-12559.php> (last access: 4 July 2013), 2013.
- Viglione, A., Chirico, G. B., Komma, J., Woods, R., Borga, M., and Blöschl, G.: Quantifying space–time dynamics of flood event types, *J. Hydrol.*, 394, 213–229, 2010.
- Viglione, A., Merz, R., Salinas, J. S., and Blöschl, G.: Flood frequency hydrology: 3. A Bayesian analysis, *Water Resour. Res.*, 49, 675–692, 2013.
- Wardekker, J. A., de Jong, A., Knoop, J. M., and van der Sluijs, J. P.: Operationalising a resilience approach to adapting an urban delta to uncertain climate changes, *Technol. Forecast. Soc.*, 77, 987–998, 2010.

---

## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

---

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[⏪](#)[⏩](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)



## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion




**Table A1.** Catchment area and mean elevation of the catchments used in this paper (Fig. 1).

Stream gauge	Stream	Catchment area (km <sup>2</sup> )	Mean elevation (m a.s.l.)
Achleiten	Danube	76 650	829
Donauwörth	Danube	15 100	640
Greimpersdorf	Ybbs	1100	676
Hofkirchen	Danube	45 610	622
Kienstock	Danube	96 000	817
Korneuburg	Danube	101 500	803
Landshut	Isar	7900	702
Obergäu	Lammer	395	1195
Oberndorf	Salzach	6120	1338
Rosenheim	Mangfall	1090	790
Schärding	Inn	25 665	1257
Schwabelweis	Danube	35 450	614
St. Johann	Kitzbüheler Ache	330	1312
Staudach	Tiroler Achen	944	1148
Steyr	Enns	5915	1138
Wasserburg	Inn	11 980	1674
Weißbach am Lofer	Saalach	567	1346
Wels	Traun	3425	819
Wildungsmauer	Danube	104 000	784
Wilhering	Danube	76 450	820

## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.



**Fig. 1.** Upper Danube basin upstream of Wildungsmauer. Red circles indicate stream gauges used in this paper. Black circles indicate the cities of Vienna and Passau. For catchment areas and mean elevations of the catchments see Table A1.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.




**Fig. 2.** Geopotential height fields (in m) at 1000 hPa of the Northern Hemisphere for latitudes above 20°. Five-day moving averages, centred on 26 May and 30 May 2013. Based on the NCEP-NCAR Reanalysis datasets (Kistler et al., 2001).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.




**Fig. 3.** Sea level pressure (hPa) in Central Europe on 31 May 2013 (00:00), 10 August 2002 (12:00), 8 July 1954 (00:00), 13 September 1899 (06:00) (all times in UTC). 2013, 2002 and 1954 are based on the NCEP-NCAR Reanalysis, while 1899 is from Lauda (1900). Circles indicate location of Passau. Times have been chosen as to be most relevant for the precipitation production.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.



**Fig. 4.** Observed precipitation totals of four large flood events in the Upper Danube basin: 29 May 00:00 to 4 June 2013 24:00; 4 August 00:00 to 15 August 2002 24:00; 7 July to 12 July 1954, and 8 September to 14 September 1899 (based on available daily records, 07:00 to 07:00. 2013 and 2002 are based on rain gauge data interpolated by the INCA method using radar (Haiden et al., 2011). 1954 and 1899 are based on rain gauge data interpolated manually within the Danube basin (HZB, 1955; Lauda, 1900). Red line indicates the Upper Danube catchment boundary above Wildungsmauer.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc


Printer-friendly Version

Interactive Discussion



## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.




**Fig. 5.** Top: catchment precipitation for the Weißbach catchment (567 km<sup>2</sup>) in May and June 2013 (dark blue shows hourly intensities, light blue cumulative precipitation). Bottom: catchment average temperatures for the Weißbach catchment (dark red) and air temperature at the Loferer Alm station (1623 m a.s.l., orange).

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)

## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.




**Fig. 6.** Cumulative precipitation (blue lines) and cumulative runoff depths (red lines, direct runoff only) for the June 2013 flood event. Light green line in the left panel shows the catchment area with temperatures below 0°C. Left: Weißbach catchment (567 km<sup>2</sup>), right: Hofkirchen catchment (45 610 km<sup>2</sup>).

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.



**Fig. 7.** Left: statistical analysis of annual maximum precipitation (48 h totals, 07:00–07:00 LT) for the Lofer rain gauge near Weißbach. Right: statistical analysis of annual maximum runoff for Weißbach (567 km<sup>2</sup>). Lines show fitted Gumbel distributions. Lofer data 1961–2013. Weißbach data 1959–2013. The 1954 and 1899 precipitation depths of 233 and 329 mm, respectively, are not included in the figure.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶

Back

Close

Full Screen / Esc

Printer-friendly Version

Interactive Discussion



## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

Title Page

Abstract

Introduction

Conclusions

References

Tables

Figures

◀

▶

◀

▶


Back

Close

Full Screen / Esc

Printer-friendly Version


Interactive Discussion



**Fig. 8.** Propagation of the June 2013 flood along the stream network of the Danube basin. Red circles indicate the stream gauges as in Fig. 1. The scale shown relates to all hydrographs.


## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.



**Fig. 9.** Top: runoff hydrographs for the June 2013 and August 2002 events at the confluence of the Inn (Schärding) and the Bavarian Danube (Hofkirchen). Runoff at Achleiten is the sum of Schärding and Hofkirchen plus the Ilz, a minor tributary. Bottom: water level hydrographs for the July 1954 and September 1899 events at similar locations (Vilshofen is close to Hofkirchen). The 1954 peak discharges at Schärding and Vilshofen were  $6300$  and  $3320 \text{ m}^3 \text{ s}^{-1}$ , respectively, and the 1899 peak discharges were  $6400$  and  $2370 \text{ m}^3 \text{ s}^{-1}$ , respectively.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)



**Fig. 10.** Travel times of the 2013, 2002, 1954 and 1899 floods along the Austrian reach of the Danube from Achleiten near Passau to Wildungsmauer near the Slovak–Austrian border. 2002-1 and 2002-2 relate to the first and second flood waves of the 2002 event, respectively. At Passau/Achleiten the flood peaks occurred on 3 June 2013, 18:00 LT; 8 August 2002, 02:00 LT; 13 August 2002, 12:00 LT; 10 July 1954, 08:00 LT; 15 September 1899, 10:00 LT, which has been plotted as 0 for the five floods in the figure.

## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.

[Title Page](#)

[Abstract](#)

[Introduction](#)

[Conclusions](#)

[References](#)

[Tables](#)

[Figures](#)

[⏪](#)

[⏩](#)

[⏴](#)

[⏵](#)

[Back](#)

[Close](#)

[Full Screen / Esc](#)


[Printer-friendly Version](#)

[Interactive Discussion](#)



## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.




**Fig. 11.** Statistical analysis of annual maximum runoff for Kienstock (95 970 km<sup>2</sup>, left) and Korneuburg (101 536 km<sup>2</sup>, right). Lines show fitted Gumbel distributions. Kienstock data 1893–2013, Korneuburg data 1828–2013. Kienstock is a combined series with the nearby Stein-Krems gauge; Korneuburg is a combined series with the nearby Vienna gauge.

[Title Page](#)
[Abstract](#)
[Introduction](#)
[Conclusions](#)
[References](#)
[Tables](#)
[Figures](#)
[⏪](#)
[⏩](#)
[◀](#)
[▶](#)
[Back](#)
[Close](#)
[Full Screen / Esc](#)
[Printer-friendly Version](#)
[Interactive Discussion](#)


## The June 2013 flood in the Upper Danube basin

G. Blöschl et al.



**Fig. 12.** Comparison of the 2013 flood (dark colours) and the 2012 flood (light colours) for the Kienstock and Korneuburg stream gauges at the Danube which are 73 km apart.

[Title Page](#)[Abstract](#)[Introduction](#)[Conclusions](#)[References](#)[Tables](#)[Figures](#)[◀](#)[▶](#)[◀](#)[▶](#)[Back](#)[Close](#)[Full Screen / Esc](#)[Printer-friendly Version](#)[Interactive Discussion](#)