

[Datei](#) [Bearbeiten](#) [Ansicht](#) [Chronik](#) [Lesezeichen](#) [Extras](#) [Hilfe](#)

Publikationsdatenbank x EW1 2015 x October 2014 : Nature Clim... x

www.nature.com/nclimats/archive/issue.html?year=2014&month=10

Meistbesucht [www.kreditkarte.at](#) [Erste Schritte](#) [Aktuelle Nachrichten](#) [Outlook Web App](#) <http://eeg.tuwien.ac.at/>

[nature.com](#) | [Publications A-Z index](#) | [Browse by subject](#) [Login](#) | [Register](#) | [Cart](#)

nature plants

First Issue
Now Online

nature climate change

Search [Go](#)

[Advanced search](#)

[Home](#) | [Current issue](#) | [News & comment](#) | [Research](#) | [Archive](#) | [Authors & referees](#) | [About the journal](#)

Archive

Archive

[2007](#) [2008](#) [2009](#) [2010](#) [2011](#) [2012](#) [2013](#) [2014](#) [2015](#)

Nature Climate Change
October 2014

[Editorials\(2\)](#) | [Correspondence\(9\)](#) | [Commentaries\(2\)](#) | [Interview](#) | [Market Watch](#) | [Research](#)
[Highlights\(4\)](#) | [News and Views\(3\)](#) | [Perspectives\(3\)](#) | [Letters\(5\)](#) | [Articles\(3\)](#) | [Corrigendum](#)

Editorials

Science that matters
 Climate change research can influence policy decisions, but needs to stretch its boundaries.
 See also: [Commentary by Sabine Fuss et al.](#)

What's your story?
 An understanding of the personal frameworks we use to absorb and contextualise climate change

nature plants

FROM BENCH
TO BIOSPHERE

Submit your research today

natureJobs.com

[Science jobs](#) | [Science events](#)

Ten doctoral positions at the International
 Max Planck Research School (IMPRS) for
 Neural Circuits
 Max Planck Institute for Brain Research

Post-Doctoral Position
 Leibniz University of Hannover

http://www.nature.com/nplants/info/gta/TWT_mc_id:BAN_NaturePlants_1406_Submit250

11:26
12.01.2015

Betting on negative emissions

Sabine Fuss, Josep G. Canadell, Glen P. Peters, Massimo Tavoni, Robbie M. Andrew, Philippe Ciais, Robert B. Jackson, Chris D. Jones, Florian Kraxner, Nebosja Nakicenovic, Corinne Le Quéré, Michael R. Raupach, Ayyoob Sharifi, Pete Smith and Yoshiki Yamagata

Bioenergy with carbon capture and storage could be used to remove CO₂ from the atmosphere, thereby achieving ‘negative emissions’. But its credibility as a climate-change mitigation option is unproven and its widespread deployment in climate stabilization scenarios might become a dangerous distraction.

Future warming will depend strongly on the cumulative CO₂ emissions released through to the end of this century^{1,2}. A finite quota of CO₂ emissions, no more than 1,200 GtCO₂, is needed from 2015 onwards in order to stabilize climate below a global average of 2°C above pre-industrial conditions by 2100 with a likelihood of 66%. This corresponds to about 30 years at current emissions levels³. However, during the past decade, emissions from fossil fuel combustion and cement production have increased substantially to 36.1 ± 1.8 Gt CO₂/year in 2013, projected to reach 37.0 (34.8-39.3) Gt CO₂/year in 2014^{3,4,5}, 65% above their 1990 level. Staying within the 2°C limit in a cost-effective way will require strong mitigation action across all sectors, with greater effort needed the further mitigation is delayed.

Actions that could stabilize climate as desired include the deliberate removal of CO₂ from the atmosphere by human intervention – called here ‘negative emissions’. Along with afforestation, the production of sustainable bioenergy with carbon capture and storage (BECCS) is explicitly being put forth as an important mitigation option by a majority of Integrated Assessment Models’ (IAMs) scenarios aiming at keeping warming below 2°C in the IPCC’s Fifth Assessment Report (AR5)⁶. Indeed, in these scenarios, IAMs often foresee absorption of CO₂ via BECCS up to (and in some cases exceeding) 1,000 GtCO₂ cumulative over the course of the century⁷, effectively doubling the available carbon quota.

BECCS is the negative emissions technology most widely selected by IAMs to meet the requirements of temperature limits of 2°C and below. It is based on assumed carbon-neutral bioenergy (i.e. the same amount of CO₂ is sequestered at steady state by biomass feedstock growth as is released during energy generation), combined with capture of CO₂ produced by combustion and its subsequent storage in geological or ocean repositories. In other words, BECCS is a net transfer of CO₂ from the atmosphere, through the biosphere, into geological layers, providing in addition a non-fossil

fuel source of energy. Other options include afforestation, direct air capture, and increases in soil carbon storage. Afforestation and increased soil carbon storage differ from BECCS in that these land use and management changes are associated with a saturation of CO₂-removal over time, and in that the sequestration is reversible with terrestrial carbon stocks inherently vulnerable to disturbance⁸.

The need for negative emissions

The IPCC's Working Group 3 (WG3) considered in AR5 over 1,000 emission pathways to 2100 (Fig. 1a). Most scenarios (101 of 116) leading to concentration levels of 430–480 ppm CO₂-equivalent (CO₂eq), consistent with limiting warming below 2°C, require global net negative emissions in the second half of this century, as do many scenarios (235 of 653) that reach between 480 and 720 ppm CO₂eq in 2100 (Fig. 1b, scenarios below zero). About half of the scenarios feature BECCS exceeding 5% of primary energy supply. Many of those (252 of 581) have net positive emissions in 2100 (Fig. 1b). Thus, BECCS does not ensure net negative emissions (i.e., its use need not completely offset all positive emissions). BECCS is an important mitigation technology especially as the stabilization level is lowered, and if near-term mitigation is delayed. By eventually requiring deeper emissions reductions, BECCS can help reconciling higher interim CO₂-eq concentrations with low long-term stabilization targets, particularly if overshooting of concentrations is allowed. Taking into account the full scenario range, global net negative emissions would need to set in around 2070 for the most challenging scenarios and progressively later for higher-temperature stabilization levels.

IAMs⁶ and Earth system models (ESMs²) provide different but complementary approaches for quantifying negative emissions requirements. ESMs simulate the compatible net CO₂ emissions based on mass balance between atmospheric changes in CO₂ and land and ocean carbon sinks. A model intercomparison of 10 ESMs found that two-thirds of the models required net negative emissions in the second half of the century⁹, but the ESMs make no assumption on how this is technically achieved. For IAMs, negative emissions are an outcome of an economic optimization driven by a choice between reducing emissions and BECCS (gross negative emissions). Both approaches model the link between CO₂ emissions, atmospheric concentrations and subsequent climate change. Importantly, some of the non-CO₂ emissions (e.g., CH₄ and N₂O from agriculture) will be very difficult to mitigate completely, as will some CO₂ emissions from industry and transportation below which mitigation will be economically and technically very difficult¹⁰. Therefore, to reach long-term climate stabilization under 2°C, there is likely to be a requirement for gross negative CO₂ emissions (i.e. at the project level) and likely also for net negative emissions (i.e. the global net balance).

The challenges ahead

The deployment of large-scale bioenergy faces biophysical, technical, and social challenges¹¹, and CCS is yet to be implemented widely. Four major uncertainties need to be resolved: (1) the physical constraints on BECCS, including sustainability of large-scale deployment relative to other land and biomass needs, such as food security and biodiversity conservation, and the presence of safe, long-term storage capacity for carbon; (2) the response of natural land and ocean carbon sinks to negative emissions; (3) the costs and financing of an untested technology; and (4) socio-institutional barriers such as public acceptance of new technologies and the related deployment policies. In the IAM scenarios in AR5⁵ that are consistent with warming of less than 2°C, the requirement for BECCS ranges between 2–10 GtCO₂ annually in 2050, corresponding to 5–25% of 2010 CO₂ emissions and 4–22% of baseline 2050 CO₂ emissions. Huge upscaling efforts will be needed to reach this level. By comparison, the current global mean removal of CO₂ by the ocean and terrestrial carbon sinks is 9.2±1.8 GtCO₂ and 10.3±2.9 GtCO₂, respectively^{5,12}. Concerning the capture and storage portion of the BECCS chain, the International Energy Agency's CCS roadmap clearly illustrates that huge efforts would be needed to achieve the scale of CCS (both fossil fuel emissions CCS and BECCS) foreseen in current stabilization scenarios, as publicly supported demonstration programs are still struggling to deliver actual large-scale projects¹³.

It is difficult to estimate the actual costs of BECCS, as it is partially in competition for resources (land, biomass, and storage capacity and cost of CCS) used in other mitigation options and for objectives beyond climate stabilization. However, while negative emissions might appear more expensive than established mitigation options including fossil fuel emissions CCS, the mitigation pathways to 2100 excluding negative emission technologies are all substantially more expensive than the pathways including those technologies^{6,14,15}.

Policymakers will need a much more complete picture of negative emissions than what is currently at hand. Issues of governance and behavioral transformations need to be better understood. The reliance of current scenarios on negative emissions, despite very limited knowledge, calls for a major new transdisciplinary research agenda to (1) examine consistent narratives for the potential of implementing and managing negative emissions, (2) estimate uncertainties and feedbacks within the socio-institutional, techno-economic and Earth system dimensions, and (3) offer guidance on how to act under the remaining uncertainties. Similarly, technological and institutional roadmaps, and rapid implementation of pilot projects are needed to test feasibility and understand the barriers to technological development.

In addition to characterizing the potential for negative emissions more reliably and geographically explicitly^{16,17}, the tradeoffs related with the use of negative emissions need to be further assessed.

Some recent collaborative modeling efforts have provided important insights on such potential tradeoffs (e.g. ref. 18). In the case of BECCS, tradeoffs are associated with (1) competition for land and possible conflicts with the objectives for food security, biodiversity conservation and the demand for water resources in different sectors (e.g. ref. 19), and (2) the existence of sufficient potential for secure and accessible storage of captured CO₂ in competition with fossil fuel CCS, uncertainties about the possibility to upscale negative emissions technologies quickly, and public acceptance.

A consistent narrative of negative emissions management therefore has four components (Fig. 2) relating to the key uncertainties. The first component refers to technological aspects: with BECCS being the negative emissions technology most widely applied by IAMs, the implied heavy demands for sustainable biomass availability are suggested to be at least 100 and up to more than 300 EJ per year of equivalent primary energy by 2050²⁰. Also, CO₂ storage potential in geological layers (aquifers, depleted fossil carbon reservoirs) and other resources such as water and fertilizer in the face of increasing food demand will need to be addressed. Bioenergy and water recycling with solar-powered distillation, algae grown offshore and fertilized with previously captured CO₂, and other innovations are among possible technologies enabling negative emissions to be achieved with lower pressure on land biomass production. However, such technologies require significant new research and development.

The second component in Fig. 2 describes carbon cycle uncertainties and dynamics in the Earth system. If negative emissions options such as BECCS are used only after significant climate change, then the response of the global carbon cycle can make the necessary amount of negative emissions even larger than for a scenario where the future CO₂ trajectory is contained below 430–480 ppm. This could occur through decreasing terrestrial and ocean sink efficiencies due to climate change, and net releases of CO₂ by the land and ocean reservoirs due to CO₂ removal over several decades^{6,12,21,22,6}.

The third component acknowledges that negative emissions will be part of a wider mitigation effort and their deployment will depend on the cost, risks and timing profile of other options. The spectrum ranges from more established mitigation technologies – for which it might then be too late – to solar radiation management geo-engineering options, which are quicker and cheaper to ramp up, but which embody a much larger scale of mostly unknown risks²³ and are not able to deal with other consequences of increased CO₂ concentrations such as the ocean acidification. This emphasizes that we are not in a position to discard the negative emissions option easily despite the above challenges. The fact that negative emissions solutions like BECCS will require time to achieve sufficient scale confirms that the future option space depends strongly on today's decisions.

The final component is concerned with institutional and policy challenges. CO₂ removal will be expensive and contentious, whereas emissions will remain cheap in the absence of strong climate policies. Therefore, any CO₂ removal strategy requires an extraordinary global regulatory framework taking into account national economic conditions. In the absence of a global climate agreement requiring stringent mitigation efforts and given the asymmetric distribution of mitigation potentials, negative emissions could help to offset emissions from countries that might not participate in reduction efforts or have less capacity to do so. This could open new perspectives on global climate management. Rigorous monitoring, reporting and verification will be needed to facilitate these options.

The development of consistent negative emission narratives is not a call for large-scale BECCS deployment, but a call to carefully and quickly assess all dimensions of its use for climate stabilization. Determining how safe it is to bet on negative emissions in the second half of this century to avoid dangerous climate change should be among our top priorities.

Acknowledgements

This work is a collaborative effort under the MaGNET (**M**anaging **G**lobal **N**egative **E**missions **T**echnologies) initiative of the Global Carbon Project (www.globalcarbonproject.org), a joint project of IGBP, IHDP, WCRP and Diversitas. J.G.C. thanks the support of the Australian Climate Change Science Program. C.D.J. was supported by the Joint UK DECC/Defra Met Office Hadley Centre Climate Programme (GA01101). G.P.P. and R.M.A. thank the support of the Norwegian Research Council (236296). Y.Y. and A.S. acknowledge that GCP Tsukuba office activities are supported by CGER/NIES. F.K. acknowledges support from IIASA's Tropical Flagship Initiative (TFI). M.T. acknowledges the Italian Ministry of Education, University and Research and the Italian Ministry of Environment, Land and Sea under the GEMINA project. C.L.Q. thanks the support of UK's Natural Environment Research Council (NE/103002X/1). R.B.J. acknowledges the U.S. Department of Agriculture (AFRI #2012-00857).

Contributions

All authors contributed to the planning of the paper. S.F. led the work together with J.G.C. and prepared Figure 2 including description of the framework benefitting from discussions with all authors. G.P.P., R.M.A. and M.T. prepared Figure 1 and/or provided the associated analysis. All authors contributed to writing the paper, providing comments to the framework and input in terms of numbers and references backing the analysis.

Sabine Fuss^{1,2*}, Josep G. Canadell³, Glen P. Peters⁴, Massimo Tavoni⁵, Robbie M. Andrew⁴, Philippe Ciais⁶, Robert B. Jackson⁷, Chris D. Jones⁸, Florian Kraxner², Nebosja Nakicenovic², Corinne Le Quéré⁹, Michael R. Raupach¹⁰, Ayyoob Sharifi¹¹, Pete Smith¹², Yoshiki Yamagata¹¹

¹Mercator Research Institute on Global Commons and Climate Change, Working Group Resources and International Trade, Torgauer Strasse 12–15 Berlin, 10829, Germany

²International Institute for Applied Systems Analysis, Schlossplatz 1, 2361 Laxenburg, Austria

³CSIRO Oceans and Atmospheric Flagship, GPO Box 3023, Canberra, ACT 2601, Australia

⁴Center for International Climate and Environmental Research - Oslo (CICERO), Gaustadalléen 21, 0349, Oslo, Norway

⁵Fondazione Eni Enrico Mattei and Centro-Mediterraneo sui Cambiamenti Climatici (CMCC) and Politecnico di Milano, Corso Magenta 63, Milan, Italy

⁶Laboratoire des Sciences du Climat et de l'Environnement, Centre d'Etudes de Orme des Merisiers - BAT 709, Gif sur Yvette 91191, France

⁷Stanford University, School of Earth Sciences, Woods Institute for the Environment, and Precourt Institute for Energy, 473 Vio Ortega, Stanford 94305, USA

⁸Met Office Hadley Centre, FitzRoy Road, Exeter, EX1 3PB, U.K. ⁹Tyndall Centre for Climate Change Research, University of East Anglia, Norwich NR4 7TJ, UK

¹⁰Climate Change Institute, Fenner School, Australian National University, Canberra, ACT 0200, Australia

¹¹National Institute for Environmental Studies, Onogawa 16-2, Tsukuba Ibaraki, 305-8506, Japan

¹²Institute of Biological & Environmental Sciences, University of Aberdeen, 23 St Machar Drive, Aberdeen, AB24 3UU, UK

*e-mail: fuss@mcc-berlin.net

References

1. Allen, M. R. *et al.* *Nature* **458**, 1163–1166 (2009).
2. *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (IPCC, 2013) [eds Stocker, T.F. *et al.*] (Cambridge University Press, Cambridge, UK and NY, 2013).
3. Friedlingstein P. *et al.* *Nature Geoscience* (in press, 2014).
4. Boden, T. A. *et al.* *Global, Regional, and National Fossil-Fuel CO₂ Emissions*. (Oak Ridge National Laboratory, U.S. Department of Energy, Oak Ridge, Tennessee, USA, 2013).
5. Le Quéré, C. *et al.* *Earth Syst. Sci. Data* **6**, 235–263. (2014).
6. Clarke, L. *et al.* In: *Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds Edenhofer, O. *et al.*) (IPCC, 2014) Cambridge University Press, Cambridge, UK and NY (accepted but not approved in detail by the 12th Session of Working Group III and the 39th Session of the IPCC on 12 April 2014 in Berlin, Germany).
7. Tavoni, M. & Socolow, R. *Climatic Change* **118**, 1–14. (2013).
8. Raupach M. R. & Canadell, J. G. in *The Continental-Scale Greenhouse Gas Balance of Europe* (eds Dolman A. J. *et al.*) (Springer, New York, (2008).
9. Jones, C. *et al.* *J. Climate* **26**, 4398–4413 (2013).
10. Davis, S.J., Caldeira, K., Matthews, H.D. *Science* **29**, 1330-1333 (2010).
11. Creutzig, F. *et al.* *Global Change Biology* (in press, 2014).
12. Ciais, P. *et al.* In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (eds Stocker, T.F. *et al.*). (IPCC, 2013) Cambridge University Press, Cambridge, UK and NY.

13. Scott, V., Gilfillan, S., Markusson, N., Chalmers, H., Haszeldine, R.S. *Nature Clim. Change* **3**, 105–111 (2012).
14. Fuss, S., Reuter, W.-H., Szolgayova, J, Obersteiner, M. *Climatic Change* **118**, 73–87. (2013).
15. Kriegler, E., Edenhofer, O., Reuster, L., Luderer, G., Klein, D. *Climatic Change* **118**, 45–57 (2013).
16. Kraxner, F. *et al. Rene* **61**, 102–108 (2014).
17. Kato, E. & Yamagata, Y. *Earth's Future* (in press, 2014).
18. Popp, A. *et al. Climatic Change* **123**, 495–509 (2014).
19. *Chapter 20 - Land and Water: Linkages to Bioenergy*. In: *Global Energy Assessment - Toward a Sustainable Future* (GEA, 2012). Cambridge University Press, Cambridge, UK and NY, International Institute for Applied Systems Analysis, Laxenburg, Austria, pp. 1459-1526.
20. Smith, P. *et al.* In: *Climate Change 2014: Mitigation of Climate Change. Contribution of Working Group III to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (IPCC, 2014) [eds Edenhofer, O. *et al.*] (Cambridge University Press, Cambridge, UK and NY (accepted but not approved in detail by the 12th Session of Working Group III and the 39th Session of the IPCC on 12 April 2014 in Berlin, Germany).
21. Cao, L. & Caldeira, K. *Environ Res. Lett.* **5**, 024011 (6pp) (2010).
22. Vichi, M., Navarra, A., Fogli, P.G. *Climatic Change* **118**, 105–118. (2013).
23. Kravitz, B. *et al. J. Geophys. Res. Atmos.* **118**, 8320–8332. (2013).
24. *Summary for Policymakers*. In: *Climate Change 2013: The Physical Science Basis. Contribution of Working Group I to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change* (IPCC, 2013) [eds Stocker, T. F. *et al.*] (Cambridge University Press, Cambridge, UK and NY, 2013).




Figure 1: CO₂ emission pathways until 2100 and the extent of net negative emissions and BECCS in 2100. Historical emissions from fossil-fuel combustion and industry (black) are primarily from the Carbon Dioxide Information Analysis Center^{4,6}. They are compared with the IPCC- AR5-WG3 emissions scenarios (pale colours) and to the four Representative Concentration Pathways (RCPs) used to project climate change in the IPCC-AR5-WG1 (dark colours) (a). The emission scenarios have been grouped into climate categories⁵ measured in ppm CO₂-equivalent in 2100 (legend) from all components and linked to the most relevant RCP. The temperature increase (right of Fig. 1a) refers to the warming in the late 21st century (2081 –2100 average) relative to the 1850–1900 average²⁴. Only scenarios assigned to climate categories are shown (1,089 of 1,184). Most scenarios that keep climate warming below 2°C use BECCS and many require net negative emissions (i.e. BECCS exceeding fossil fuel emissions) in 2100 (b). Data source: AR5 database and GCP/CDIAC.




Figure 2: Four components of consistent negative emission narratives.