# AN INVOLUTION ON DYCK PATHS THAT PRESERVES THE RISE COMPOSITION AND INTERCHANGES THE NUMBER OF RETURNS AND THE POSITION OF THE FIRST DOUBLE FALL

#### MARTIN RUBEY

ABSTRACT. Motivated by a recent paper of Adin, Bagno and Roichman, we present an involution on Dyck paths that preserves the rise composition and interchanges the number of returns and the position of the first double fall.

### 1. INTRODUCTION

Let  $\mathfrak{S}_n(321)$  be the set of 321-avoiding permutations on n letters. In a recent paper of Adin, Bagno and Roichman [1] a new equidistribution result of two statistics on  $\mathfrak{S}_n(321)$ is used to demonstrate the Schur positivity of quasi-symmetric functions associated with certain subsets of  $\mathfrak{S}_n(321)$ . Using a recursively defined bijection<sup>1</sup> the authors establish

$$\sum_{\mathfrak{S}_n(321)} \mathbf{x}^{\mathrm{LRMAX}(\pi)} q^{\mathrm{bl}(\pi)} = \sum_{\pi \in \mathfrak{S}_n(321)} \mathbf{x}^{\mathrm{LRMAX}(\pi)} q^{n-\mathrm{ldes}(\pi^{-1})}, \text{ where }$$

- LRMAX( $\pi$ ) is the set of positions of the left-to-right maxima of  $\pi$  and  $\mathbf{x}^{L} = \prod_{i \in L} x_{i}$ ,
- $bl(\pi) = |\{i : \forall j \le i : \pi(j) \le i\}|$  is the number of blocks of  $\pi$  and
- $\operatorname{ldes}(\pi) = \max(\{0\} \cup \{i : \pi(i) > \pi(i+1)\})$  is the position of the last descent of  $\pi$ .

We exhibit an involution which proves the stronger result that, fixing LRMAX, the joint distribution of  $\pi \mapsto bl(\pi)$  and  $\pi \mapsto n - ldes(\pi^{-1})$  is in fact symmetric.

Let us first transform the statistics involved into statistics on Dyck paths, using a bijection due to Krattenthaler [3]<sup>2</sup>. To do so, it is convenient to define a Dyck path as a path in  $\mathbb{N}^2$  consisting of (1,0)- and (0,1)-steps beginning at the origin, ending at (n,n), and never going below the diagonal y = x. We refer to (1,0)-steps also as *east steps* or *falls*, and to (0,1)-steps also as *north steps* or *rises*, and the number n as the *semilength* of the path. Krattenthaler's bijection maps

- $\mathfrak{S}_n(321)$  to the set of Dyck paths  $\mathfrak{D}_n$  of semilength n,
- LRMAX( $\pi$ ) to RISES(D), the set of x-coordinates (plus 1) of the north steps of the Dyck path,
- $bl(\pi)$  to ret(D), the number of *returns* of the path, that is, the number of east steps that end on the main diagonal,
- $\operatorname{ldes}(\pi^{-1})$  to  $\operatorname{ldr}(D)$ , the *y*-coordinate of the midpoint of the last double rise, that is, the last pair of two consecutive north steps. For the path without double rises, we set  $\operatorname{ldr}(D) = 0$ .

 $\pi \in 0$ 

Supported by the Austrian Science Fund (FWF): P 29275.

<sup>&</sup>lt;sup>1</sup>http://www.findstat.org/Mp00124

<sup>&</sup>lt;sup>2</sup>http://www.findstat.org/Mp00119

#### MARTIN RUBEY

The purpose of this note is to provide an involution  $\Phi^3$  that implies

$$\sum_{\pi \in \mathfrak{D}_n} \mathbf{x}^{\mathrm{RISES}(\pi)} p^{\mathrm{ret}(D)} q^{n-\mathrm{ldr}(D)} = \sum_{\pi \in \mathfrak{D}_n} \mathbf{x}^{\mathrm{RISES}(\pi)} p^{n-\mathrm{ldr}(D)} q^{\mathrm{ret}(D)}.$$

### 2. An involution on Dyck paths

In the spirit of  $[2]^4$ , we first introduce an invertible map  $\phi$  taking a Dyck path D that ends with two east steps and satisfies ldr(D) < n-1 (that is, not maximal) to a Dyck path  $\phi(D)$  that ends with two east steps and has at least two returns. The crucial properties of this map are as follows:

- RISES $(\phi(D)) = \text{RISES}(D)$
- $\operatorname{ret}(\phi(D)) = \operatorname{ret}(D) + 1$
- $\operatorname{ldr}(\phi(D)) = \operatorname{ldr}(D) + 1$

The desired involution  $\Phi$  then maps an arbitrary path  $D(NE)^{\ell}$  (of semilength  $n + \ell$ ) to  $\phi^{(n-\operatorname{ldr}(D)-\operatorname{ret}(D))}(D)$   $(NE)^{\ell}$ .

To define  $\phi$ , write D as the concatenation of three paths P, Q and R where

- R is maximal such that it has no double rises, begins with a rise and the path Q before R ends with an east step, and
- QR is a prime Dyck path, that is, its only return is the final east step.

Let  $N_0 Q_0 N_1 Q_1 \dots N_k Q_k$  be the prime decomposition of Q, that is, all the  $Q_i$  are prime Dyck paths and the  $N_i$  are sequences of north steps. Note that  $N_0$  is non-empty, because Q does not return to the diagonal. We distinguish three cases, illustrated in Figure 1:

(1) If k = 0 or  $N_1 \neq \emptyset$  define  $\phi(D) = P Q_0 N_1 \dots N_k Q_k N_0 R$ .

In the remaining two cases let N be all but one of the north steps of the final rise in Q, and let Q' be Q with  $N_0$ ,  $Q_0$  and N erased.

- (2) If  $N_1 = \cdots = N_k = \emptyset$  define  $\phi(D) = P N Q_0 Q' N_0 R$ .
- (3) Otherwise define  $\phi(D) = P Q_0 N Q' N_0 R$ .

We now provide an inverse for  $\phi$ . Let D be a Dyck path that ends with two east steps and has at least two returns. Write  $\tilde{D}$  as the concatenation of paths  $P, \tilde{Q}, \tilde{E}, \tilde{N}$ and R where

- R is maximal such that it has no double rises and begins with a rise, and
- $\tilde{Q} \,\tilde{E} \,\tilde{N} \,R$  is a Dyck path with precisely two returns,  $\tilde{N}$  is a maximal sequence of north steps and  $\tilde{E}$  is a maximal sequence of east steps.

Let  $\tilde{N}_0 \tilde{Q}_1 \tilde{N}_1 \dots \tilde{Q}_\ell \tilde{N}_\ell$  be the prime decomposition of  $\tilde{Q}$ .

- (1) If  $\tilde{Q}$  ends with a double rise, we have  $\phi^{(-1)}(\tilde{D}) = P \tilde{N} \tilde{Q} \tilde{E} R$ . (2) Otherwise, if  $\tilde{Q}$  has no returns,  $\phi^{(-1)}(\tilde{D}) = P \tilde{N} \tilde{Q}_1 \tilde{N}_1 \dots \tilde{Q}_\ell \tilde{N}_\ell \tilde{N}_0 \tilde{E} R$ .
- (3) Otherwise,  $\phi^{(-1)}(\tilde{D}) = P \tilde{N} \tilde{Q}_1 \tilde{Q}_2 \tilde{N}_2 \dots \tilde{Q}_\ell \tilde{N}_\ell \tilde{N}_1 \tilde{E} R.$

To conclude, we show that applying the constructions in Cases (1), (2) and (3) above a path corresponding to Cases (1), (2) and (3), respectively. Suppose that  $D = \phi(D)$  is obtained by applying Case (1). Because  $Q_k$  ends with the last double rise of D followed by east steps, Q ends with a double rise.

<sup>&</sup>lt;sup>3</sup>http://www.findstat.org/Mp00132

<sup>&</sup>lt;sup>4</sup>http://www.findstat.org/Mp00118



FIGURE 1. Examples for Dyck paths and their images under  $\phi$ 

In Case (2),  $N Q_0 Q'$  is a prime Dyck path whose last sequence of north steps has length 1 so  $\tilde{Q}$  does not end with a double rise and has no returns, whereas  $\tilde{Q}\tilde{E}$  is a prime Dyck path.

Finally, in Case (3),  $NQ'N_0$  has no returns and its last sequence of north steps has length 1. Thus,  $\tilde{Q}$  does not end with a double rise, but does have a return coming from  $Q_0$ .

### References

- R. M. Adin, E. Bagno and Y. Roichman, Block decomposition of permutations and Schur-positivity, J. Algebraic Combin. (to appear); arχiv:1611.06979.
- S. Elizalde and M. Rubey, Symmetries of statistics on lattice paths between two boundaries, Adv. Math. 287 (2016), 347–388.
- [3] C. Krattenthaler, Permutations with restricted patterns and Dyck paths, Adv. Appl. Math. 27 (2001), 510–530.

# MARTIN RUBEY

*E-mail address*: Martin.Rubey@tuwien.ac.at

FAKULTÄT FÜR MATHEMATIK UND GEOINFORMATION, TU WIEN, AUSTRIA